

Literature

- Andersson, C. Risebrobakken, B, Jansen, E, and Dahl, SO. 2003. Late Holocene surface ocean conditions of the Norwegian Sea (Voring Plateau). *Paleoceanography*, 18: Art. no. 1044
- Andrus, C.F.T., Crowe, D.E., Sandweiss, D.H., Reitz, E.J. and Romanek, C.S. 2002. Otolith $\delta^{18}\text{O}$ record of mid-Holocene sea surface temperatures in Peru. *Science*, 295:1508-1511.
- Berg E. and Pedersen T. (2001). Variability in recruitment, growth and sexual maturity of coastal cod (*Gadus morhua* L.) in a fjord system in northern Norway. *Fisheries Research*, 52:179-189
- Campana, S.E. 1999. Chemistry and composition of fish otoliths: pathways, mechanisms and applications. *Marine Ecology Progress Series*, 188:263-297.
- Campana, S.E. and Thorrold, S.R. 2001. Otoliths, increments, and elements: key to a comprehensive understanding of fish populations. *Canadian Journal of Fisheries and Aquatic Sciences*, 58:30-38.
- Carpenter, S.J., Erickson J.M. and Holland Jr, F.D. 2003. Migration of a Late Cretaceous fish. *Nature*, 423: 70-74.
- Dettman, D.L. and Lohmann, K.C. 1995. Microsampling carbonates for stable isotope and minor element analysis: physical separation of samples on a 20 micrometer scale. *Journal of Sedimentary Research A, Sedimentary Petrology and Processes*, 65:566-569.
- Folkvord, A., Rukan, K., Johannessen, A. and Moksness, E. 1997. Early life history of herring larvae in contrasting feeding environments determined by otolith microstructure analysis. *Journal of Fish Biology*, 51:250-263.
- Geffen, A.J., Pearce, N.J. and Perkins, W.P. 1998. Metal concentrations in fish otoliths in relation to body composition after laboratory exposure to mercury and lead. *Marine Ecology Progress Series*, 165:235-245
- Geffen, A.J., Jarvis, K., Thorpe, J.P., Leah, R.T. and Nash, R.D.M. 2003. Trace element concentrations in the otoliths of plaice *Pleuronectes platessa* and whiting *Merlangius merlangus* in the eastern Irish Sea. *Netherlands Journal of Sea Research*, 50: 245-254
- Høie, H., Andersson, C., Folkvord, A. and Karlsen, Ø. 2004. Precision and accuracy of stable isotope signals in otoliths of pen reared cod (*Gadus morhua* L.) when sampled with high resolution micromill. *Marine Biology*, 144:1039-1049.
- Ivany, L.C., Patterson, W.P. and Lohmann, K. 2000. Cooler winters as a possible cause of mass extinctions at the Eocene/Oligocene boundary. *Nature*, 407:887-890.
- Köster, F.W., Schnack, D. and Möllmann, C. 2003. Scientific knowledge of biological processes that are potentially useful in fish stock predictions. *Scientia Marina*, 67:101-127.
- Løken, S. and Pedersen, T. (1996). Effect of parent type and temperature on vertebrae number in juvenile cod, *Gadus morhua* (L.), in Northern Norway. *Sarsia*, 80: 293-298.
- Løken, S., Pedersen, T., and Berg, E. (1994). Vertebrae numbers as an indicator for recruitment mechanism of coastal cod of northern Norway. *ICES Marine Science Symposium*, 198: 510-519.
- Møller, D. (1968). Genetic differences between cod groups in the Lofoten area. *Nature*, 212:824.
- Ottersen, G., Planque, B., Belgrano, A., Post, E., Reid, P.C. and Stenseth, N.C. 2001. Ecological effects of the North Atlantic oscillation. *Oecologia*, 128:1-14.
- Patterson, W.P. 1998. North American continental seasonality during the last millennium: high-resolution analysis of sagittal otoliths. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 138:271-303.
- Patterson, W.P. 1999. Oldest isotopically characterized fish otoliths provide insight to Jurassic continental climate of Europe. *Geology*, 27:199-202.
- Pogson, G.H. and Fevolden, S.E. (2003). Natural selection and the genetic differentiation of coastal and Arctic populations of the Atlantic cod in northern Norway: a test involving nucleotide sequence variation at the pantophysin (*PanI*) locus. *Molecular Ecology*, 12: 63-74.
- Rollefsen, G. (1934). The cod otolith as a guide to race, sexual development and mortality. *Rapports et Procès-verbaux des Réunions. Conseil international pour l'Exploration de la Mer*, 88: 1-6.
- Thresher, R. E. (1999). Elemental composition of otoliths as a stock delineator in fishes. *Fisheries Research*, 43: 165-204.
- Svåsand, T., Jørstad, K.E., Otterå, H., and Kjesbu, O.S. (1996). Differences in growth performance between Arcto-Norwegian and Norwegian coastal cod reared under identical conditions. *Journal of Fish Biology*, 49: 108-119.
- Van Heel M., Harauz G., Orlova E.V., Schmidt R. & Schatz M. 1996. A new generation of the IMAGIC image processing system. *Journal of Structural Biology*, 116:17-24.
- Wurster, C.M. and Patterson, W.P. 2001. Late Holocene climate change for the eastern interior United States: evidence from high-resolution $\delta^{18}\text{O}$ values of sagittal otoliths. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 170:81-100.
- Wurster, C.M., Patterson, W.P. and Cheatham, M.M. 1999. Advances in microdrilling techniques: a new apparatus for acquiring high-resolution oxygen and carbon stable isotope values and major/minor elemental ratios from accretionary carbonate. *Computers & Geosciences*, 25:1159-1166.
- Øiestad, V. 1994. Historic changes in cod stocks and cod fisheries: Northeast Arctic cod. *ICES Marine Science Symposia*, 198:17-30.