

Selfconsistent field

Interaction energy of two charges depends on their distance $|\vec{r}_1 - \vec{r}_2|$:

$$W(|\vec{r}_1 - \vec{r}_2|) = \frac{q_1 q_2}{|\vec{r}_1 - \vec{r}_2|}$$

The two charges are an electron and a little volume dV at \vec{r}_2 containing charge cloud of density ρ

$$q_1 \rightarrow (-e) \quad q_2 \rightarrow \rho(\vec{r}_2)dV \quad \rightarrow \quad \rho(\vec{r}_2)d^3 r_2$$

and the interaction energy of these two charges is

$$dW(|\vec{r}_1 - \vec{r}_2|) = \frac{(-e)\rho(\vec{r}_2)}{|\vec{r}_1 - \vec{r}_2|} d^3 r_2$$

and summing over all the small volume elements means integrating over the whole volume of the cloud gives the potential energy due to the interaction with a cloud

$$W(\vec{r}) = \int \frac{(-e)\rho(\vec{x})}{|\vec{r} - \vec{x}|} d^3 x$$

If the charge cloud represents one electron in state $\psi_i(\vec{x})$

$$\rho(\vec{x}) = (-e)|\psi_i(\vec{x})|^2$$

If we have N electrons, each in its state, the total density becomes

$$\rho(\vec{x}) = (-e) \sum_{i=1}^N |\psi_i(\vec{x})|^2$$

and again integrating over the whole volume of the cloud gives the potential energy due to the interaction with a (probability based density) cloud of electrons

$$W(\vec{r}) = \int \frac{(-e)^2 \sum_{i=1}^N |\psi_i(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3 x$$

Now solving the Schrödinger equation with $W(\vec{r})$,

$$(T + V + W) \psi_i(\vec{x}) = E_i \psi_i(\vec{x})$$

We first need to know the $W(\vec{r})$, but that depends on all the other N solutions

$$W(\vec{r}) = \int \frac{(-e)^2 \sum_{i=1}^N |\psi_i(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3 x$$

We thus start an approximation chain. First we choose some simple approximation, e.g. the hydrogen-like states, or we might know the states for another atom. We call it

$$\psi_i^{(0)}(\vec{x})$$

From the set of all N $\psi_i^{(0)}$ we construct

$$W^{(1)}(\vec{r}) = \int \frac{e^2 \sum_{i=1}^N |\psi_i^{(0)}(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3 x$$

In atomic units the whole Schrödinger equation is

$$\left(-\frac{1}{2}\nabla^2 - \frac{Z}{r} + W^{(1)}(\vec{r})\right) \psi_i^{(1)}(\vec{x}) = E_i^{(1)} \psi_i^{(1)}(\vec{x})$$

Now we show the chain in steps: first step

$$\begin{aligned} \psi_i^{(0)}(\vec{x}) &\rightarrow W^{(1)}(\vec{r}) = \int \frac{\sum_{i=1}^N |\psi_i^{(0)}(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3x \\ \left(-\frac{1}{2}\nabla^2 - \frac{Z}{r} + W^{(1)}(\vec{r})\right) \psi_i^{(1)}(\vec{x}) &= E_i^{(1)} \psi_i^{(1)}(\vec{x}) \end{aligned}$$

Second step

$$\begin{aligned} \psi_i^{(1)}(\vec{x}) &\rightarrow W^{(2)}(\vec{r}) = \int \frac{\sum_{i=1}^N |\psi_i^{(1)}(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3x \\ \left(-\frac{1}{2}\nabla^2 - \frac{Z}{r} + W^{(2)}(\vec{r})\right) \psi_i^{(2)}(\vec{x}) &= E_i^{(2)} \psi_i^{(2)}(\vec{x}) \end{aligned}$$

Third step

$$\begin{aligned} \psi_i^{(2)}(\vec{x}) &\rightarrow W^{(3)}(\vec{r}) = \int \frac{\sum_{i=1}^N |\psi_i^{(2)}(\vec{x})|^2}{|\vec{r} - \vec{x}|} d^3x \\ \left(-\frac{1}{2}\nabla^2 - \frac{Z}{r} + W^{(3)}(\vec{r})\right) \psi_i^{(3)}(\vec{x}) &= E_i^{(3)} \psi_i^{(3)}(\vec{x}) \end{aligned}$$

This chain can continue, until the set of $\psi_i^{(n)}$ produces a potential $W^{(n+1)}$ which is the same as $W^{(n)}$, which was the one to determine $\psi_i^{(n)}$. The potentials and functions become consistent, hence the name *Selfconsistent field*.

Criterion for self-consistency:

$$\int \sum_{i=1}^N \left| |\psi_i^{(2)}(\vec{x})|^2 - |\psi_i^{(1)}(\vec{x})|^2 \right| d^3x < \epsilon$$

where $\epsilon \propto 10^{-8}$

The total energy of N electrons and connection to Hartree-Fock theory

Please, correct and fill in the text as exercise

Is the energy of N-electrons approximately equal to the sum of energies from the above solutions?

$$\sum_{i=1}^N (E_i) \approx E?$$

$$\left[\sum_{i=1}^N \left(-\frac{1}{2}\nabla_{r_i}^2 - \frac{Z}{r_i} \right) + \sum_{(i,j) \text{ pairs}} \frac{1}{|\vec{r} - \vec{x}|} \right] \Phi(r_1, r_2, \dots, r_N) = E \Phi(r_1, r_2, \dots, r_N)$$

$$\Phi(r_1, r_2, \dots, r_N) \approx \psi_1(r_1)\psi_2(r_2)\dots\psi_N(r_N)$$

$$\sum_{i=1}^N (E_i) \approx E?$$

$$\sum_{j=1}^N (E_j) \propto \sum_{j=1}^N \int \psi_j^*(\vec{y}) \left(\int \frac{e^2 \sum_{i=1}^N |\psi_i(\vec{x})|^2 d^3x}{|\vec{y} - \vec{x}|} \right) \psi_j(\vec{y}) d^3y$$

$$\sum_{(i,j) \text{ pairs}} \int \psi_j^*(\vec{y}) \left(\int \frac{e^2 \sum_{i=1}^N |\psi_i(\vec{x})|^2 d^3x}{|\vec{y} - \vec{x}|} \right) \psi_j(\vec{y}) d^3y$$

$$\sum_{(i,j) \text{ pairs}} f(r_j, r_i) = \frac{1}{2} \sum_{j=1}^N \sum_{i=1, i \neq j}^N f(r_j, r_i)$$

$$E \approx \sum_{i=1}^N (E_i) - \sum_{(i,j) \text{ pairs}} \int \psi_j^*(\vec{y}) \left(\int \frac{e^2 \sum_{i=1}^N |\psi_i(\vec{x})|^2 d^3x}{|\vec{y} - \vec{x}|} \right) \psi_j(\vec{y}) d^3y$$

$$\Phi_{a,b,\dots,N}^{HF}(r_1, r_2, \dots, r_N) \rightarrow \left| \begin{array}{l} \psi_a(r_1)\psi_b(r_1)\dots\psi_N(r_1) \\ \psi_a(r_2)\psi_b(r_2)\dots\psi_N(r_2) \\ \dots\dots\dots \\ \psi_a(r_N)\psi_b(r_N)\dots\psi_N(r_N) \end{array} \right|$$

$$\left\langle \psi_a \psi_b \left| \frac{1}{|r-\bar{x}|} \right| \psi_a \psi_b \right\rangle \rightarrow \left\langle \psi_a \psi_b \left| \frac{1}{|r-\bar{x}|} \right| \psi_a \psi_b \right\rangle - \left\langle \psi_a \psi_b \left| \frac{1}{|r-\bar{x}|} \right| \psi_b \psi_a \right\rangle$$

$$\langle a, b | f(r_1, r_2) | a, b \rangle \rightarrow \langle a, b | f(r_1, r_2) | a, b \rangle - \langle a, b | f(r_1, r_2) | b, a \rangle$$