

Infinite types that satisfy the principle of omniscience

Martín Hötzel Escardó

University of Birmingham, UK

Bergen, Norway, 21th January 2016

Mathematics in dependent type theory

1. I'll work in intensional Martin-Löf type theory (MLTT).
2. I will make a number of remarks related to HoTT, in particular regarding -1 -truncation and equivalence.
3. Sometimes I will use *function extensionality*.

Alternatively, I can assume that our hypothetical functions are extensional in a suitable sense, like Bishop did. However, this leads to *setoid hell*.

4. I will work informally but rigorously.

But I have also written formal versions of the proofs in Agda notation.

LPO

For any given $p : \mathbb{N} \rightarrow 2$, we can either find $n : \mathbb{N}$ with $p(n) = 0$, or else determine that $p(n) = 1$ for all $n : \mathbb{N}$.

$$\Pi(p : \mathbb{N} \rightarrow 2).(\Sigma(n : \mathbb{N}).p(n) = 0) + (\Pi(n : \mathbb{N}).p(n) = 1)$$

LPO

For any given $p : \mathbb{N} \rightarrow 2$, we can either find $n : \mathbb{N}$ with $p(n) = 0$, or else determine that $p(n) = 1$ for all $n : \mathbb{N}$.

$$\Pi(p : \mathbb{N} \rightarrow 2).(\Sigma(n : \mathbb{N}).p(n) = 0) + (\Pi(n : \mathbb{N}).p(n) = 1)$$

For any given $p : \mathbb{N} \rightarrow 2$, we can either find a root of p , or else determine that there is none.

$$\Pi(p : \mathbb{N} \rightarrow 2).(\Sigma(n : \mathbb{N}).p(n) = 0) + \neg(\Sigma(n : \mathbb{N}).p(n) = 0)$$

Subsingleton version of LPO

Any $p : \mathbb{N} \rightarrow 2$ either has a root or it doesn't.

$$\Pi(p : \mathbb{N} \rightarrow 2). \|\Sigma(n : \mathbb{N}). p(n) = 0\| + \neg(\Sigma(n : \mathbb{N}). p(n) = 0)$$

(No need to -1 -truncate the rightmost Σ , as the negation of a type is equivalent to the negation of its truncation.)

The LPO types

$$\prod(p : \mathbb{N} \rightarrow 2).(\sum(n : \mathbb{N}).p(n) = 0) + \neg(\sum(n : \mathbb{N}).p(n) = 0)$$

and

$$\prod(p : \mathbb{N} \rightarrow 2).\|\sum(n : \mathbb{N}).p(n) = 0\| + \neg(\sum(n : \mathbb{N}).p(n) = 0)$$

are logically (but not necessarily homotopically) equivalent.

The LPO types

$$\prod(p : \mathbb{N} \rightarrow 2).(\sum(n : \mathbb{N}).p(n) = 0) + \neg(\sum(n : \mathbb{N}).p(n) = 0)$$

and

$$\prod(p : \mathbb{N} \rightarrow 2).\|\sum(n : \mathbb{N}).p(n) = 0\| + \neg(\sum(n : \mathbb{N}).p(n) = 0)$$

are logically (but not necessarily homotopically) equivalent.

The second is a retract of the first.

This doesn't use the HoTT formulation of the axiom of choice.

It is an instance of choice that just holds.

LPO is undecided

$$\prod(p : \mathbb{N} \rightarrow 2).(\sum(n : \mathbb{N}).p(n) = 0) + (\neg \sum(n : \mathbb{N}).p(n) = 0)$$

1. A meta-theorem is that MLTT doesn't inhabit LPO or \neg LPO.
2. Each of them is consistent with MLTT.

Classical models validate LPO.

Effective and continuous models validate \neg LPO.

3. LPO is undecided, and we'll keep it that way.
4. But we'll say it is a *taboo*.

We now make \mathbb{N} larger by adding a point at infinity

Let \mathbb{N}_∞ be the type of decreasing binary sequences.

$$\mathbb{N}_\infty \stackrel{\text{def}}{=} \Sigma(\alpha : 2^{\mathbb{N}}). \Pi(n : \mathbb{N}). \alpha(n) = 0 \rightarrow \alpha(n + 1) = 0.$$

1. As you know, \mathbb{N} is the *initial algebra* of the functor $1 + (-)$.
2. \mathbb{N}_∞ is the *final coalgebra* of this functor.

(This requires function extensionality.)

We now make \mathbb{N} larger by adding a point at infinity

Let \mathbb{N}_∞ be the type of decreasing binary sequences.

$$\mathbb{N}_\infty \stackrel{\text{def}}{=} \Sigma(\alpha : 2^{\mathbb{N}}). \Pi(n : \mathbb{N}). \alpha(n) = 0 \rightarrow \alpha(n+1) = 0.$$

1. The type \mathbb{N} embeds into \mathbb{N}_∞ by mapping the number $n : \mathbb{N}$ to the sequence $\underline{n} \stackrel{\text{def}}{=} 1^n 0^\omega$.
2. A point not in the image of this is $\infty \stackrel{\text{def}}{=} 1^\omega$.
3. The assertion that every point of \mathbb{N}_∞ is of one of these two forms is equivalent to LPO.
4. What is true is that no point of \mathbb{N}_∞ is different from all points of these two forms.
5. The embedding $\mathbb{N} + 1 \rightarrow \mathbb{N}_\infty$ is an isomorphism iff LPO holds.
6. But the complement of its image is empty. We say it is **dense**.

Theorem

$$\prod(p : \mathbb{N}_\infty \rightarrow 2).(\sum(x : \mathbb{N}_\infty).p(x) = 0) + \neg \sum(x : \mathbb{N}_\infty).p(x) = 0$$

1. This is LPO with \mathbb{N} replaced by \mathbb{N}_∞ .
2. We don't use continuity axioms, which anyway are not available in MLTT.
3. However, this is motivated by topological (not homotopical) considerations.

In Johnstone's *topological topos*, \mathbb{N}_∞ gets interpreted as the one-point compactification of discrete \mathbb{N} .

Here we are seeing a *logical manifestation of topological compactness*.

4. This theorem actually makes sense in any variety of constructive mathematics (JSL 2013).

WLPO is also undecided by MLTT

$$\prod(p : \mathbb{N} \rightarrow 2).(\prod(n : \mathbb{N}).p(n) = 1) + \neg \prod(x : \mathbb{N}).p(x) = 1$$

(This implies that every Turing machine carries on for ever or it doesn't.)

But we have:

Theorem $\prod(p : \mathbb{N}_\infty \rightarrow 2).(\prod(n : \mathbb{N}).p(\underline{n}) = 1) + \neg \prod(n : \mathbb{N}).p(\underline{n}) = 1$

1. The point is that now we quantify over \mathbb{N} , although the function p is defined on \mathbb{N}_∞ .
2. This again holds in any variety of constructive mathematics and doesn't rely on continuity axioms (JSL'2013).

Some consequences

1. Every function $f : \mathbb{N}_\infty \rightarrow \mathbb{N}$ is constant or not.
2. Any two functions $f, g : \mathbb{N}_\infty \rightarrow \mathbb{N}$ are equal or not.
3. Any function $f : \mathbb{N}_\infty \rightarrow \mathbb{N}$ has a minimum value, and it is possible to find the point at which the minimum value is attained.
4. For any function $f : \mathbb{N}_\infty \rightarrow \mathbb{N}$ there is a point $x : \mathbb{N}_\infty$ such that if f has a maximum value, the maximum value is x .
5. Any function $f : \mathbb{N}_\infty \rightarrow \mathbb{N}$ is not continuous, or not-not continuous.
6. There is a non-continuous function $f : \mathbb{N}_\infty \rightarrow \mathbb{N}$ iff WLPO holds.

Are there more types like \mathbb{N}_∞ ?

1. Plenty.
2. Our business here is how to construct them.
3. But we pause to reflect first.

What have we been doing?

Giving examples of types X and properties P of X such that the assertion

for all $x : X$, either $P(x)$ or not $P(x)$

just holds.

1. In classical mathematics, we assume excluded middle.
2. Here we investigate how much of it just holds.

No philosophy or meta-mathematics, except for side-remarks.

We just prove mathematical theorems.

Two notions

Definition (Omniscient type)

A type X is **omniscient** if for every $p : X \rightarrow 2$, the assertion that we can find $x : X$ with $p(x) = 0$ is decidable.

In symbols:

$$\prod (p : X \rightarrow 2). (\sum (x : X). p(x) = 0) + (\neg \sum (x : X). p(x) = 0).$$

Two notions

Definition (Omniscient type)

A type X is **omniscient** if for every $p : X \rightarrow 2$, the assertion that we can find $x : X$ with $p(x) = 0$ is decidable.

In symbols:

$$\Pi(p : X \rightarrow 2).(\Sigma(x : X).p(x) = 0) + (\neg\Sigma(x : X).p(x) = 0).$$

Definition (Searchable type)

A type X is **searchable** if for every $p : X \rightarrow 2$ we can find $x_0 : X$, called a *universal witness* for p , such that if $p(x_0) = 1$, then $p(x) = 1$ for all $x : X$.

In symbols,

$$\Pi(p : X \rightarrow 2).\Sigma(x_0 : X).p(x_0) = 1 \rightarrow \Pi(x : X).p(x) = 1.$$

Their relationship

$$\text{omniscient}(X) \stackrel{\text{def}}{=} \Pi(p : X \rightarrow 2).(\Sigma(x : X).p(x) = 0) + (\neg \Sigma(x : X).p(x) = 0).$$

$$\text{searchable}(X) \stackrel{\text{def}}{=} \Pi(p : X \rightarrow 2).\Sigma(x_0 : X).p(x_0) = 1 \rightarrow \Pi(x : X).p(x) = 1.$$

NB. These types are not subsingletons in general.

Proposition A type X is searchable iff it has a point and is omniscient:

$$\text{searchable}(X) \iff X \times \text{omniscient}(X).$$

A few theorems rely on pointedness, using the notion of searchability.

Closure under Σ

If X is omniscient/searchable and Y is an X -indexed family of omniscient/searchable types, then so is its disjoint sum $\Sigma(x : X).Y(x)$.

Closure under Π

Not to be expected in general.

E.g. \mathbb{N}_∞ and 2 are omniscient, but in continuous and effective models of type theory, the function space $\mathbb{N}_\infty \rightarrow 2$ is not.

In the topological topos, $\mathbb{N}_\infty \rightarrow 2$ is a countable discrete space.

Closure under finite products

Theorem A product of searchable types indexed by a finite type is searchable.

Brouwerian closure under countable products

Theorem Brouwerian intuitionistic axioms \implies

A countable product of searchable types is searchable.

This is a kind of Tychonoff theorem, if we think of searchability as a “synthetic” notion of compactness.

In particular, the Cantor type $2^{\mathbb{N}}$, which is interpreted as the Cantor space in the topological topos, is searchable.

Brouwerian closure under countable products

Theorem Brouwerian intuitionistic axioms \implies

A countable product of searchable types is searchable.

This is a kind of Tychonoff theorem, if we think of searchability as a “synthetic” notion of compactness.

In particular, the Cantor type $2^{\mathbb{N}}$, which is interpreted as the Cantor space in the topological topos, is searchable.

1. Falsified in one effective model
(the effective topos, which is realizability over Kleene's K_1).

Brouwerian closure under countable products

Theorem Brouwerian intuitionistic axioms \implies

A countable product of searchable types is searchable.

This is a kind of Tychonoff theorem, if we think of searchability as a “synthetic” notion of compactness.

In particular, the Cantor type $2^{\mathbb{N}}$, which is interpreted as the Cantor space in the topological topos, is searchable.

1. Falsified in one effective model
(the effective topos, which is realizability over Kleene's K_1).
2. But validated in another effective model
(realizability over Kleene's K_2),
and in the topological topos.

(I implemented this in Agda, by disabling the termination checker in a particular function. One can run interesting examples.)

We will need this form of closure under Π

Theorem (micro Tychonoff)

A product of searchable types indexed by a subsingleton type is itself searchable.

That is, if X is a subsingleton, and Y is an X -indexed family of searchable types, then the type $\Pi(x : X).Y(x)$ is searchable.

We will need this form of closure under Π

Theorem (micro Tychonoff)

A product of searchable types indexed by a subsingleton type is itself searchable.

That is, if X is a subsingleton, and Y is an X -indexed family of searchable types, then the type $\Pi(x : X).Y(x)$ is searchable.

I don't think this can be proved if searchability is replaced by omniscience (that is, if we don't assume that every $Y(x)$ is pointed).

This is easy with excluded middle, but we are not assuming it.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0)$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0)$.
6. $(X \rightarrow 0) \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0)$.
6. $(X \rightarrow 0) \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).(X \rightarrow 0) \rightarrow p(z) = 1$.

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0)$.
6. $(X \rightarrow 0) \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.
 $p(z_0) = 1 \rightarrow \prod(z : Z).(X \rightarrow 0) \rightarrow p(z) = 1$.
7. By transitivity of \rightarrow , we get

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.
2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.
4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.
5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

$$p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1.$$

$$p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0).$$

6. $(X \rightarrow 0) \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

$$p(z_0) = 1 \rightarrow \prod(z : Z).(X \rightarrow 0) \rightarrow p(z) = 1.$$

7. By transitivity of \rightarrow , we get

$$p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow p(z) = 1, \text{ so}$$

Theorem A subsingleton-indexed product of searchable types is searchable.

1. Let X subsingleton, $Y(x)$ searchable for every $x : X$.

2. $Z \stackrel{\text{def}}{=} \prod(x : X).Y(x)$.

We have $\prod(x : X).(Z \simeq Y(x))$ and $(X \rightarrow 0) \rightarrow (Z \simeq 1)$.

3. Let $p : Z \rightarrow 2$.

4. Construct $z_0(x) \stackrel{\text{def}}{=} \dots$ in Z using the first equivalence.

5. $X \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

$p(z_0) = 1 \rightarrow \prod(z : Z).X \rightarrow p(z) = 1$.

$p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow (X \rightarrow 0)$.

6. $(X \rightarrow 0) \rightarrow p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$.

$p(z_0) = 1 \rightarrow \prod(z : Z).(X \rightarrow 0) \rightarrow p(z) = 1$.

7. By transitivity of \rightarrow , we get

$p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 0 \rightarrow p(z) = 1$, so

$p(z_0) = 1 \rightarrow \prod(z : Z).p(z) = 1$. **Q.E.D.**

Disjoint sum with a point at infinity

Theorem

The disjoint sum of a countable family of searchable sets with a point at infinity is searchable.

We need to say how we add a point at infinity.

The type $1 + \Sigma(n : \mathbb{N}).X(n)$ won't do, of course.

We will do this in a couple of steps.

Injectivity of the universe of types

Theorem

For any embedding $e : A \rightarrow B$, every $X : A \rightarrow U$ extends to some $Y : B \rightarrow U$ along e , up to equivalence,

$$\prod (a : A). (Y(e(a)) \simeq X(a)).$$

Recall that $e : A \rightarrow B$ is an embedding iff its fibers $e^{-1}(b)$,

$$\Sigma (a : A). f(a) = b,$$

are all subsingletons.

Injectivity of the universe of types

Theorem

For any embedding $e : A \rightarrow B$, every $X : A \rightarrow U$ extends to some $Y : B \rightarrow U$ along e , up to equivalence.

Two constructions:

1. We have the “maximal” extension $Y = X/e$.

$$\begin{aligned}(X/e)(b) &= \Pi (s : e^{-1}(b)) . X(\text{pr}_1 s) \\ &\simeq \Pi (a : A) . e(a) = b \rightarrow X(a).\end{aligned}$$

Injectivity of the universe of types

Theorem

For any embedding $e : A \rightarrow B$, every $X : A \rightarrow U$ extends to some $Y : B \rightarrow U$ along e , up to equivalence.

Two constructions:

1. We have the “maximal” extension $Y = X/e$.

$$\begin{aligned}(X/e)(b) &= \Pi (s : e^{-1}(b)) . X(\text{pr}_1 s) \\ &\simeq \Pi (a : A) . e(a) = b \rightarrow X(a).\end{aligned}$$

2. And also the “minimal” extension $Y = X \setminus e$.

$$\begin{aligned}(X \setminus e)(b) &= \Sigma (s : e^{-1}(b)) . X(\text{pr}_1 s) \\ &\simeq \Sigma (a : A) . e(a) = b \times X(a).\end{aligned}$$

The first one works our purposes.

Injectivity of the universe of types

Let $e : A \rightarrow B$ be an embedding and $X : A \rightarrow U$.

Consider the extended type family $X/e : B \rightarrow U$ defined above:

$$(X \setminus e)(b) = \Pi (s : e^{-1}(b)) . X(\text{pr}_1 s)$$

We have

1. For all $b : B$ not in the image of the embedding,

$$(X/e)(b) \simeq 1.$$

2. If for all $a : A$, the type $X(a)$ is searchable too, then for all $b : B$ the type $(X/e)(b)$ is searchable, by **micro-Tychonoff**.
3. Hence if additionally B is searchable, the type $\Sigma(b : B).(X/e)(b)$ is searchable too.
4. We are interested in $A = \mathbb{N}$ and $B = \mathbb{N}_\infty$, which gives the disjoint sum of $X(a)$ with a point at infinity.

A map $L : (\mathbb{N} \rightarrow U) \rightarrow U$

Let $e : \mathbb{N} \rightarrow \mathbb{N}_\infty$ be the natural embedding.

Given $X : \mathbb{N} \rightarrow U$, first take $X/e : \mathbb{N}_\infty \rightarrow U$

This step adds a point at infinity to the sequence.

We then sum over \mathbb{N}_∞ , to get $L(X)$:

$$L(X) = \Sigma(u : \mathbb{N}_\infty).(X/e)(u).$$

Then L maps any sequence of searchable types to a searchable type.

Iterating this map $L : (\mathbb{N} \rightarrow U) \rightarrow U$

We get (very large!) searchable ordinals, with the property that any inhabited *decidable* subset has a least element.

Iterating this map $L : (\mathbb{N} \rightarrow U) \rightarrow U$

We get (very large!) searchable ordinals, with the property that any inhabited *decidable* subset has a least element.

They are all countable.

Or rather they each have a countable subset with empty complement.

Iterating this map $L : (\mathbb{N} \rightarrow U) \rightarrow U$

We get (very large!) searchable ordinals, with the property that any inhabited *decidable* subset has a least element.

They are all countable.

Or rather they each have a countable subset with empty complement.

An ordinal is a type X with a transitive, extensional, accessible relation $(-) < (-) : X \rightarrow X \rightarrow U$.

Iterating this map $L : (\mathbb{N} \rightarrow U) \rightarrow U$

We get (very large!) searchable ordinals, with the property that any inhabited *decidable* subset has a least element.

They are all countable.

Or rather they each have a countable subset with empty complement.

An ordinal is a type X with a transitive, extensional, accessible relation $(-) < (-) : X \rightarrow X \rightarrow U$.

1. **Extensional** means that any two elements with the same predecessors are equal.
2. The **accessibility** of points of X is inductively defined.

We say that $x : X$ is accessible whenever every $y < x$ is accessible.

The accessibility of a point is a subingleton.

3. $<$ is accessible if every $x : X$ is accessible.

The accessibility of $<$ implies that it is subsingleton valued, and that X is set.

A functor $F : U \rightarrow U$

$F(X) = L(\lambda n.X)$, which is equivalent to $\Sigma(u : \mathbb{N}_\infty).\Pi(n : \mathbb{N}).X^{e(n)=u}$.

An equivalent coninductive definition of F is given by constructors

zero : $X \rightarrow F(X)$,
succ : $F(X) \rightarrow F(X)$.

1. The Cantor type $2^{\mathbb{N}}$ is the carrier of a final coalgebra of F .
2. There is an initial algebra, whose carrier is the subset of Cantor consisting of the sequences with finitely many zeros, for a suitable notion of finiteness.

(Which is classically equivalent to the classical one.)

End